III. Enumerate the following:

(30 marks)

- 1) 3 (three) opportunistic infections occurring in AIDS.
- 2) 3 (three) causes of haemorrhage.
- 3) 3 (three) complications of intestinal bilharziasis.
- 4) 3 (three) viral diseases affecting the nervous system.
- 5) 3 (three) endogenous pigments.
- 6) 3 (three) causes of ischaemia.
- 7) 3 (three) sites of primary complex.
- 8) 3 (three) skeletal manifestations of rickets.
- 9) 3 (three) aetiological factors involved in the development of cancer (carcinogenesis).
- 10) 3 (three) types of acute suppurative inflammation.

IV. Give an account on:

(30 marks)

- 1. Causes of cell injury.
- 2. Complications of tuberculosis of the vertebrae (Pott's disease).
- 3. Morphologic features (types) of candidiasis (moniliasis).
- 4. Methods of spread of malignant tumours.
- 5. Manifestations of secondary acquired syphilis.
- 6. Effects of portal hypertension.

End of the exam
With best wishes

Faculty of Nursing 2nd Semester Final Exam Time allowed: 2 Hours (29/5/2016)

2-	All	are	adverse	effect	of	thiazide	diuretic	except:
----	-----	-----	---------	--------	----	----------	----------	---------

(Hyponatremia - Hypokalemia - Hypocalcemia).

3- One of the following is a common indication of nifedipine:

(Hypertension - Tachycardia - Tinnitus).

4- The following is one of insulin uses:

(Hypoglycemia - Hypersensitivity - Hyperkalemia).

5- If aspirin used for treatment of fever in children due to viral infection, it leads to

(Renal Damage - Reye's Syndrome - Metabolic Acidosis).

III- Mention True OR False: (5 Marks)

- 1. Tolerance is a common side effect of nitrate if used for a long period.
- 2. It is safe to stop Na⁺ nitroprusside infusion suddenly after control of emergency hypertension.
- 3. Protamine sulphate is the specific antidote for warfarin toxicity.
- 4. Barbiturates used for treatment of neonatal hyperbilirubinemia.
- 5. Metformin is a hypoglycemic drug used to treat thin type II diabetic patient.

GOOD LUCK

Dr. Mohamed Balaha.

4- Herbal medicine:

(Safe – Mask Manifestation of Disease – Not Interact With Other Drugs).

5- Which of the following drugs used in treatment of acute hypotension?

(Dobutamine – Noradrenaline - Fenoldopam).

III- Mention True OR False: (10 Marks)

- 1. Plants is not one of drugs sources.
- 2. Probenecid inhibits renal tubular secretion of penicillin.
- 3. Isoprenaline used in treatment of acute bronchial asthma.
- 4. Allergy is a predictable pharmacological response of drug.
- 5. Large dose of aspirin not follow zero order kinetic.
- 6. Lipophilic β -Blocker is the drug of choice in essential familial tremors.
- 7. Dobutamine is selective D_1 receptor agonist.
- 8. Initial syncopal attack is one of side effects of prazosin.
- 9. Acetylcholine used effectively clinically.
- 10. Oily and insoluble substances can be administer by intravenous route of administration.

GOOD LUCK

Dr. Mehamed Balaha